

Optical geometry analysis of the electromagnetic self-force

Sebastiano Sonego* and Marek A. Abramowicz^{†‡}

**Università di Udine, Via delle Scienze 208, 33100 Udine, Italy*

†Department of Physics, Chalmers University of Technology, 41296 Göteborg, Sweden

‡Nicolaus Copernicus Astronomical Centre, Polish Academy of Sciences, Warsaw, Poland

16 February 2006; L^AT_EX-ed February 7, 2008

Abstract

We present an analysis of the behaviour of the electromagnetic self-force for charged particles in a conformally static spacetime, interpreting the results with the help of optical geometry. Some conditions for the vanishing of the local terms in the self-force are derived and discussed.

PACS: 04.20.Cv, 04.70.-s, 04.90.+e

Keywords: Optical geometry; self-force; radiation reaction

*sebastiano.sonego@uniud.it

†marek@fy.chalmers.se

1 Introduction

The motion of a point particle with mass m , electric charge e , and four-velocity¹ v^a , is described by the equation

$$ma_a = K_a + F_a, \quad (1.1)$$

where $a_a := v^b \nabla_b v_a$ is the particle four-acceleration, K_a is the sum of the external four-forces acting on the particle, and F_a accounts for the back-reaction of the particle own electromagnetic field on its motion. In contrast to a well-established tradition, we shall avoid referring to F_a as the *radiation reaction*, because such a terminology is somehow improper and can be misleading [4, 5]; we shall use the term *self-force* instead. According to the classic analysis of DeWitt-Brehme-Hobbs [2, 3], F_a can be written as the sum of two contributions, $F_a = F_a^{(l)} + F_a^{(nl)}$, where²

$$F_a^{(l)} = \frac{2}{3} e^2 k_a{}^b v^c \nabla_c a_b + \frac{1}{3} e^2 k_a{}^b R_{bc} v^c \quad (1.2)$$

and

$$F_a^{(nl)} = e^2 v^b \int_{-\infty}^{\tau} d\tau' f_{aba'} v^{a'}. \quad (1.3)$$

Here v^a and τ are the four-velocity and proper time of the particle, $k_a{}^b = \delta_a{}^b + v_a v^b$ is the projector onto the three-space orthogonal to v^a , R_{ab} is the Ricci tensor, and $f_{aba'}$ is a bi-tensor associated with the presence of “tails” in the electromagnetic field [2, 7]. The term $F_a^{(l)}$ is the relativistic generalisation to a curved spacetime of the non-covariant expression $(2/3)e^2 \ddot{\mathbf{v}}$ for the self-force [4, 5], and can be regarded as purely local, as it depends only on quantities evaluated at the actual position of the particle. On the contrary, the contribution $F_a^{(nl)}$ depends on the entire past history of the particle and on the property of spacetime of being able to back-scatter electromagnetic waves [2, 7], encapsulated in the quantity $f_{aba'}$, and represents therefore an essentially non-local contribution. The two terms on the right hand side of (1.2) are known as the von Laue [4, 5] and the Hobbs [3]

¹We use latin indices $a, b, \dots$ from the beginning of the alphabet as abstract indices [1], while greek letters $\mu, \nu, \dots$ denote components in some chart and run from 0 to 3. We work in units in which $G = c = 1$, and choose the positive signature for the metric. We follow the conventions of Ref. [1] for the curvature tensors; the Ricci tensor R_{ab} has then the opposite sign than in Refs. [2, 3].

²The equation of motion obtained putting together (1.1)–(1.3) contains third derivatives of the particle coordinates, hence leads to unphysical conclusions such as preacceleration and runaway solutions, as it happens for the Lorentz-Dirac equation in flat spacetime [4]. These pathologies can be removed by a reduction-of-order technique [6]. Nevertheless, in the following we shall consider the standard expression (1.2) for $F_a^{(l)}$, that differs from the one so obtained only to higher orders, and is therefore equivalent to it as long as one limits oneself to the classical domain, in which the theory is defined.

forces, respectively. We shall denote them as³

$$F_a^{(\text{vL})} := \frac{2}{3} e^2 k_a{}^b v^c \nabla_c a_b \quad (1.4)$$

and

$$F_a^{(\text{H})} := \frac{1}{3} e^2 k_a{}^b R_{bc} v^c, \quad (1.5)$$

so (1.2) can be rewritten as $F_a^{(\text{I})} = F_a^{(\text{vL})} + F_a^{(\text{H})}$. It should be clear from the outset that all this, as well as the treatment that follows, applies to classical point particles — an unrealistic model which presents pathologies, given that the size of a classical object cannot be smaller than its Compton wavelength. The extension to particles of a finite size has been discussed recently [9].

In the last decade there has been a renewed interest in calculations of self-forces of different nature. The main motivation for such a revival is the possibility to go beyond the test particle approximation when studying motion in a gravitational field, which is important for investigations about the generation of gravitational waves. (For recent reviews, see [10] and references therein.) However, due to the complexity of the calculations in the gravitational case, several authors have considered simpler models involving the scalar and electromagnetic self-interactions, which are also interesting by themselves. In particular, much attention has been paid to the non-local contribution. In this article we shall instead focus on the behaviour of the local term $F_a^{(\text{I})}$ for the electromagnetic case, arguing that it can be understood more easily in terms of the so-called optical geometry, rather than of the usual geometry of spacetime. A similar analysis could be carried out for scalar and gravitational self-forces, but we shall not cover such cases here. Also, we leave for further investigation the issue of whether optical geometry could be successfully applied to the challenging problem of computing the non-local term (see, however, the end of Sec. 5 for a brief discussion of this point).

We begin by considering, in the next section, a situation in which the charge moves uniformly on a circular orbit in a static, spherically symmetric spacetime, comparing the results in the special cases of Einstein's universe and Schwarzschild spacetime. For the latter, we shall see that $F_a^{(\text{I})} = 0$ at the closed photon orbit $r = 3M$. This property will be interpreted in Sec. 3 by introducing the notion of optical geometry, in which light paths on $t = \text{const}$ hypersurfaces are geodesics. In Sec. 4 we generalise the analysis to an arbitrary conformally static spacetime, showing that if a charge moves with constant speed along a possible light path, the self-force is the one associated with optical geometry (apart from an obvious conformal rescaling), and providing other conditions for the different parts of the self-force to vanish. Section 5 contains a brief summary of the results.

³Noteworthy, the von Laue force can also be written as

$$F_a^{(\text{vL})} = \frac{2}{3} e^2 \nabla_{(v)}^{(\text{FW})} a_a,$$

where $\nabla_{(v)}^{(\text{FW})}$ is the Fermi-Walker derivative along v^a , which for a generic field of one-forms u_a is [8]

$$\nabla_{(v)}^{(\text{FW})} u_a := v^b \nabla_b u_a + (a_a v^b - v_a a^b) u_b.$$

Hence, the von Laue term $F_a^{(\text{vL})}$ vanishes iff the acceleration a_a is Fermi-Walker transported along the particle worldline.

2 Uniform circular motion in static spherically symmetric spacetimes

Let us begin by discussing a rather special, but highly significant case. Consider a charge moving on a circular orbit $r = \text{const}$ in the plane $\theta = \pi/2$ of a static, spherically symmetric spacetime, with metric

$$g = -e^{2\Phi(r)} dt^2 + \alpha(r) dr^2 + r^2 (d\theta^2 + \sin^2 \theta d\varphi^2) , \quad (2.1)$$

where Φ and α are known functions, with α positive. The components of the four-velocity are

$$v^\mu = \Gamma (\delta_t^\mu + \Omega \delta_\varphi^\mu) , \quad (2.2)$$

where Ω is a parameter and $\Gamma = (e^{2\Phi} - \Omega^2 r^2)^{-1/2}$ is fixed by the normalization $v_\mu v^\mu = -1$. It is convenient to introduce the velocity $v = e^{-\Phi} \Omega r$ measured by static observers, and the corresponding Lorentz factor $\gamma := (1 - v^2)^{-1/2} = e^\Phi \Gamma$.

If the motion is uniform, i.e. $\Omega = \text{const}$, the only non-vanishing component of the acceleration is

$$a_r = \Gamma^2 \left(e^{2\Phi} \frac{d\Phi}{dr} - \Omega^2 r \right) , \quad (2.3)$$

and the von Laue force is given by

$$F_\mu^{(\text{vL})} = \frac{2}{3} e^2 \gamma^5 \Omega r \frac{e^{-\Phi}}{\alpha} \left(1 - r \frac{d\Phi}{dr} \right) \left(e^{2\Phi} \frac{d\Phi}{dr} - \Omega^2 r \right) (-\Omega \delta_\mu^t + \delta_\mu^\varphi) . \quad (2.4)$$

The Hobbs term is

$$F_\mu^{(\text{H})} = \frac{1}{3} e^2 \gamma^3 \Omega e^{-\Phi} (e^{-2\Phi} r^2 R_{tt} + R_{\varphi\varphi}) (-\Omega \delta_\mu^t + \delta_\mu^\varphi) , \quad (2.5)$$

where

$$R_{tt} = \frac{e^{2\Phi}}{\alpha} \left(\frac{d^2\Phi}{dr^2} + \frac{2}{r} \frac{d\Phi}{dr} + \left(\frac{d\Phi}{dr} \right)^2 - \frac{1}{2\alpha} \frac{d\Phi}{dr} \frac{d\alpha}{dr} \right) \quad (2.6)$$

and

$$R_{\varphi\varphi} = 1 - \frac{1}{\alpha} + \frac{r}{\alpha} \left(\frac{1}{2\alpha} \frac{d\alpha}{dr} - \frac{d\Phi}{dr} \right) \quad (2.7)$$

are the only relevant non-vanishing components of the Ricci tensor.⁴ Because of the rather complicated dependence on Φ , α , and their derivatives, it is more instructive to focus on two particular cases.

2.1 Einstein's universe

The metric of Einstein's static universe corresponds to $\Phi = 0$ and $\alpha(r) = (1 - r^2/R^2)^{-1}$, where R is a positive parameter (the “radius” of the three-dimensional spherical space). The coordinate $r \in [0, R[$, however, does not cover the whole manifold, so it is convenient

⁴The component $R_{\varphi\varphi}$ in (2.7) is evaluated at $\theta = \pi/2$. The general expression for $R_{\varphi\varphi}$ contains an overall extra coefficient $\sin^2 \theta$.

to introduce a new variable $\chi \in [0, \pi]$ defined through $r = R \sin \chi$. The metric then takes the form

$$g = -dt^2 + R^2 d\chi^2 + R^2 \sin^2 \chi (d\theta^2 + \sin^2 \theta d\varphi^2) , \quad (2.8)$$

showing that $R\chi$ measures the proper distance from the point $\chi = 0$. The function α becomes $\alpha(\chi) = 1/\cos^2 \chi$, and the only non-vanishing component of the acceleration is

$$a_\chi = \frac{\partial r}{\partial \chi} a_r = -\frac{\Omega^2 R^2 \sin \chi \cos \chi}{1 - \Omega^2 R^2 \sin^2 \chi} . \quad (2.9)$$

The magnitude (with sign) of the acceleration is $\gamma^2(v^2/r) \cos \chi$, so the acceleration differs from the special relativistic expression $\gamma^2 v^2/r$ only by the factor $\cos \chi$, whose origin can be easily understood by looking at Fig. 1. Note that the acceleration for orbits with




Figure 1: Embedding diagram for the surface $t = \text{const}$, $\theta = \pi/2$ in Einstein's universe, showing the geometrical meaning of the quantities R , r , and χ . The centripetal acceleration of a particle moving at $r = \text{const}$ is given by the tangential component of the acceleration in the fictitious three-dimensional Euclidean space. For $\chi < \pi/2$ the centripetal acceleration points in the direction of decreasing r , while for $\pi/2 < \chi < \pi$ it points in the direction of increasing r .

$\chi \in]\pi/2, \pi[$ has the opposite sign than for those with $\chi \in]0, \pi/2[$, and vanishes when $\chi = \pi/2$.

The von Laue force is

$$F_\mu^{(\text{vL})} = \frac{2}{3} e^2 \gamma^5 \Omega^3 R^2 \sin^2 \chi \cos^2 \chi \left(-\Omega \delta_\mu^t + \delta_\mu^\varphi \right) . \quad (2.10)$$

It also vanishes for $\chi = \pi/2$, but does not exhibit any change in sign.

Finally, let us compute the Hobbs force. The components of the Ricci tensor can be evaluated directly from the expressions (2.6) and (2.7), or simply remembering that the symmetries of Einstein's universe imply $R_{\mu\nu} = (2/R^2)h_{\mu\nu}$, where

$$\begin{aligned} h &= \left(1 - r^2/R^2\right)^{-1} dr^2 + r^2 \left(d\theta^2 + \sin^2 \theta d\varphi^2\right) \\ &= R^2 d\chi^2 + R^2 \sin^2 \chi \left(d\theta^2 + \sin^2 \theta d\varphi^2\right) \end{aligned} \quad (2.11)$$

is the metric tensor on the $t = \text{const}$ hypersurfaces [11]. Thus, we have

$$F_\mu^{(\text{H})} = \frac{2}{3} e^2 \gamma^3 \Omega \sin^2 \chi \left(-\Omega \delta_\mu^t + \delta_\mu^\varphi\right). \quad (2.12)$$

This expression is non-vanishing for all non-trivial values of χ . Therefore, the local part of the self-force,

$$F_\mu^{(\text{l})} = \frac{2}{3} e^2 \gamma^5 \Omega \sin^2 \chi \left(1 + \Omega^2 R^2 \cos 2\chi\right) \left(-\Omega \delta_\mu^t + \delta_\mu^\varphi\right), \quad (2.13)$$

never vanishes, in spite of the fact that when the motion takes place on the spatial geodesic $\chi = \pi/2$, the von Laue force does.

2.2 Schwarzschild spacetime

In Schwarzschild spacetime,

$$\Phi = \frac{1}{2} \ln \left(1 - \frac{2M}{r}\right) \quad (2.14)$$

and $\alpha = e^{-2\Phi}$. Then, from Eq. (2.3) we obtain

$$a_r = \Gamma^2 \left(\frac{M}{r^2} - \Omega^2 r\right) = \frac{M - \Omega^2 r^3}{r(r - 2M - \Omega^2 r^3)} \quad (2.15)$$

and, from (2.4),

$$F_\mu^{(\text{vL})} = \frac{2}{3} e^2 \gamma^5 \Omega r \left(1 - \frac{2M}{r}\right)^{-3/2} \left(1 - \frac{3M}{r}\right) \left(\frac{M}{r^2} - \Omega^2 r\right) \left(-\Omega \delta_\mu^t + \delta_\mu^\varphi\right). \quad (2.16)$$

Since $R_{ab} = 0$, the Hobbs force vanishes identically, so $F_\mu^{(\text{l})} = F_\mu^{(\text{vL})}$.

Let us now study the behaviour of a_r and $F_\mu^{(\text{l})}$ for different values of r . When r/M is very large (Newtonian gravity), Eq. (2.15) can be approximated as

$$a_r \approx \gamma^2 \left(\frac{M}{r^2} - \Omega^2 r\right). \quad (2.17)$$

Apart from the factor γ^2 , which represents only special-relativistic corrections, the norm of a_a is just the difference between the gravitational and the centripetal accelerations in Newtonian mechanics. This agrees with the physical interpretation of a_a , which represents the thrust per unit mass that must be applied to the particle in order to keep it at the

fixed value of r with angular velocity Ω . In particular, $a_a = 0$ for a Keplerian motion, $\Omega^2 = \Omega_K(r)^2 := M/r^3$ or $r = r_K(\Omega) := (M/\Omega^2)^{1/3}$. In the same approximation,

$$F_\mu^{(1)} \approx \frac{2}{3} e^2 \Omega r \gamma^5 \left(\frac{M}{r^2} - \Omega^2 r \right) \left(-\Omega \delta_\mu^t + \delta_\mu^\varphi \right). \quad (2.18)$$

The local part of the self-force behaves then in the following way: When the motion is Keplerian it vanishes, consistently with the fact that $a_a = 0$; in the super-Keplerian regime ($\Omega^2 > \Omega_K(r)^2$ or $r > r_K(\Omega)$) it points backward with respect to the direction of motion; finally, in the sub-Keplerian regime ($\Omega^2 < \Omega_K(r)^2$ or $r < r_K(\Omega)$) it points forward with respect to the direction of motion. In the particular case of a static charge, $\Omega = 0$ and $F_\mu^{(1)} = 0$.

It is interesting to note that this behaviour is peculiar to gravity and has no counterpart for motion in other central fields such as, e.g., a Coulomb one. This is easily seen by computing the self-force associated with circular motion in Minkowski spacetime,

$$F_\mu \equiv F_\mu^{(1)} = -\frac{2}{3} e^2 \Omega^3 r^2 \gamma^5 \left(-\Omega \delta_\mu^t + \delta_\mu^\varphi \right), \quad (2.19)$$

where the second equality follows immediately from (2.16) with $M = 0$ (see also Ref. [12], where this expression is derived explicitly for the case of synchrotron radiation). Now, the self-force always points backward with respect to the direction of motion, regardless of the magnitude of Ω . This is related to the fact that the right hand side of Eq. (2.19) does not contain any indication about the central force responsible for keeping the charge on circular motion, contrary to what happens in the gravitational case, where a term M/r^2 appears together with $\Omega^2 r$. The combination $M/r^2 - \Omega^2 r$ is the same that appears in a_r — see Eq. (2.15) —, and is ultimately linked to the association between gravity and curvature. Thus, although for most purposes one can think of Newtonian gravity as a field on a flat background, the present analysis shows that this picture would lead to incorrect conclusions as far as the self-force is concerned.⁵

Let us come back to the analysis of a_r and $F_\mu^{(1)}$. For $r > 3M$ the qualitative behaviour of these quantities at different values of r is like in the Newtonian limit examined above, even when r/M is not large. However, it is easily seen from Eqs. (2.15) and (2.16) that when $r = 3M$, $a_r = (3M)^{-1}$ and $F_\mu^{(1)} = 0$. Thus, the thrust needed in order to keep a particle on the closed photon orbit does not depend on the particle speed [13]. Furthermore, if the particle is charged, the local part of the self-force vanishes, in spite of the fact that motion takes place on a circle and is not geodesic.⁶ Finally, for orbits with $2M < r < 3M$, it follows from (2.16) that the local part of the self-force always points backward with respect to the direction of motion. Since such orbits are necessarily sub-Keplerian, as it follows from the inequality⁷

$$\left(\Omega^2 - \Omega_K(r)^2 \right) r^2 < 1 - 3M/r, \quad (2.20)$$

this behaviour is the opposite than for $r > 3M$.

⁵Apparently, this point is not appreciated in the extant literature. See, e.g., Ref. [4], p. 183, for an explicit statement that the Newtonian and Coulomb problems are alike.

⁶This point is related to earlier results [14].

⁷Proof: Consider the condition $2M/r + \Omega^2 r^2 < 1$, which guarantees that Γ be real or, equivalently, that the particle worldline be timelike; subtract $\Omega_K(r)^2 r^2$ from both sides, and use the definition $\Omega_K(r)^2 := M/r^3$.

3 Optical geometry

The behaviour of a_r can be understood in the following way. The component a_r of the acceleration, given by the expression (2.15), is proportional to the thrust necessary to keep the particle on its orbit, and can be written as the difference between a gravitational part $a_r^{(g)}$, independent of Ω , and a centripetal part $a_r^{(c)}$:

$$a_r^{(g)} = \frac{M}{r^2 (1 - 2M/r)} ; \quad (3.21)$$

$$a_r^{(c)} = \Omega^2 r \frac{r - 3M}{(1 - 2M/r)(r - 2M - \Omega^2 r^3)} . \quad (3.22)$$

(Of course, $a_r^{(g)}$ and $a_r^{(c)}$ have signs opposite to those of the gravitational and centripetal forces. Alternatively, one can identify $a_r^{(c)}$ with the centrifugal field acting on the particle in the comoving frame.) It is then evident from (3.22) that the centripetal force vanishes at $r = 3M$, for which $a_r = a_r^{(g)} = (3M)^{-1}$. In addition, (3.22) predicts that $a_r^{(c)}$ has opposite sign in the regions $r > 3M$ and $r < 3M$.

These properties admit a simple explanation if one imagines that the particle motion takes place in the so-called optical spacetime [15], with a metric $\tilde{g} = (1 - 2M/r)^{-1} g$, under the action of a “gravitational potential” Φ given by (2.14), which produces a “gravitational field” [16]

$$\tilde{g}_\mu = -k_\mu{}^\nu \nabla_\nu \Phi = - \left(1 - \frac{2M}{r}\right)^{-1} \frac{M}{r^2} \delta_\mu^r . \quad (3.23)$$

We have then $a_r^{(g)} = -\tilde{g}_r$ and, in the optical spacetime, the magnitude of the gravitational field is given by the simple Newtonian expression $(\tilde{g}^{\mu\nu} \tilde{g}_\mu \tilde{g}_\nu)^{1/2} = M/r^2$. Also, defining $\tilde{v}^\mu := (1 - 2M/r)^{1/2} v^\mu$, so that $\tilde{g}_{\mu\nu} \tilde{v}^\mu \tilde{v}^\nu = -1$, one gets

$$\tilde{a}_\mu = \tilde{g}_{\mu\sigma} \tilde{v}^\nu \tilde{\nabla}_\nu \tilde{v}^\sigma = - \frac{\Gamma^2 \Omega^2 (r - 3M)}{1 - 2M/r} \delta_\mu^r , \quad (3.24)$$

and $a_r^{(c)} = \tilde{a}_r$. The magnitude of the acceleration in optical spacetime is

$$(\tilde{g}^{\mu\nu} \tilde{a}_\mu \tilde{a}_\nu)^{1/2} = \Gamma^2 \Omega^2 |r - 3M| . \quad (3.25)$$

The presence of the factor $(r - 3M)$ in $a_r^{(c)}$ can be understood intuitively [17] considering an embedding diagram [18] of the section $\theta = \pi/2$ of the optical space $(\mathcal{S}, \tilde{h}_{ab})$, where $\mathcal{S}$ is any $t = \text{const}$ hypersurface of the Schwarzschild spacetime, and the metric $\tilde{h}_{ab}$ has the coordinate representation [15]

$$\tilde{h} = \left(1 - \frac{2M}{r}\right)^{-2} dr^2 + \left(1 - \frac{2M}{r}\right)^{-1} r^2 (d\theta^2 + \sin^2 \theta d\varphi^2) \quad (3.26)$$

(see Fig. 2 and the Appendix).

Let us now consider the self-force. The von Laue term associated with the optical quantities has the components

$$\tilde{F}_\mu^{(\text{vL})} = -\frac{2}{3} e^2 \gamma^5 \Omega^3 r^2 \left(1 - \frac{2M}{r}\right)^{-2} \left(1 - \frac{3M}{r}\right)^2 \left(-\Omega \delta_\mu^t + \delta_\mu^\varphi\right) , \quad (3.27)$$




Figure 2: Embedding diagram for part of the surface $t = \text{const}$, $\theta = \pi/2$ in Schwarzschild spacetime, with the optical geometry. The throat corresponds to $r = 3M$. The centripetal acceleration of a particle moving at $r = \text{const}$ is given by the tangential component of the acceleration in the fictitious three-dimensional Euclidean space. Notice that for $r > 3M$ the centripetal acceleration points in the direction of decreasing r , while for $2M < r < 3M$ it points in the direction of increasing r .

so it always points backward with respect to the direction of motion, vanishes only at $r = 3M$, and reduces to (2.19) in the region $r \gg M$. The Hobbs force in the optical space can be computed either directly, or by using the transformation formula (2.11) of Ref. [19]. Its components are

$$\tilde{F}_\mu^{(H)} = \frac{2}{3} e^2 \gamma^3 \Omega \frac{M}{r} \left(1 - \frac{2M}{r}\right)^{-1} \left(1 - \frac{3M}{r}\right) (-\Omega \delta_\mu^t + \delta_\mu^\varphi) . \quad (3.28)$$

For $r \gg M$, this becomes

$$\tilde{F}_\mu^{(H)} \approx \frac{2}{3} e^2 \gamma^3 \Omega \frac{M}{r} (-\Omega \delta_\mu^t + \delta_\mu^\varphi) . \quad (3.29)$$

Note that the magnitude of the von Laue force is proportional to $\Omega^3 r$, like in Minkowski space, while the magnitude of the Hobbs force is proportional to $M\Omega/r^2$, which can be rewritten as M/r^3 times the speed Ωr . Thus, from the perspective offered by optical geometry, the coefficient

$$\frac{M}{r^2} - \Omega^2 r$$

that appears in Eq. (2.18) should not be regarded as representing the difference between the gravitational and the centripetal forces, because the first term actually arises from the *curvature* of $(\mathcal{S}, \tilde{h}_{ab})$. The same coefficient is present for uniform circular motion at $r = \text{const}$ on $(\mathcal{S}, \tilde{h}_{ab})$, regardless of the nature (gravitational or other) of the “central” force. Therefore, when the local part of the self-force is analysed in optical geometry, the puzzling peculiarity of gravity that was noticed in Sec. 2.2 disappears.

It is worth pointing out that $F_\mu^{(l)}$, $\tilde{F}_\mu^{(vL)}$, and $\tilde{F}_\mu^{(H)}$ all vanish at $r = 3M$, as it follows from Eqs. (2.16), (3.27) and (3.28). However, while the property $\tilde{F}_\mu^{(vL)}(3M) = 0$ is natural from the point of view of optical geometry, because the circle $r = 3M$ is a geodesic of $(\mathcal{S}, \tilde{h}_{ab})$ and the charge motion is uniform, the fact that also $\tilde{F}_\mu^{(H)}(3M) = 0$ — hence $F_\mu^{(l)}(3M) = 0$ — is due to the algebraic properties of the optical Ricci tensor at $r = 3M$ rather than to the geodesic character in $(\mathcal{S}, \tilde{h}_{ab})$ of the closed photon orbit (see Sec. 4.2 below). Therefore, the change of sign of $F_\mu^{(l)}$ at $r = 3M$ has a different origin than for other phenomena in which a similar “reversal” takes place [20].

4 General results

If $(\mathcal{M}, g_{ab})$ and $(\mathcal{M}, \tilde{g}_{ab})$ are two conformally related spacetimes, with

$$\tilde{g}_{ab} = e^{-2\Phi} g_{ab} , \quad (4.1)$$

the local and non-local parts of the corresponding self forces are related as $F_a^{(l)} = e^{-\Phi} \tilde{F}_a^{(l)}$ and $F_a^{(nl)} = e^{-\Phi} \tilde{F}_a^{(nl)}$ [19]. Then, in order to compute $F_a^{(l)}$ or $F_a^{(nl)}$, one can work in a conformally related spacetime $(\mathcal{M}, \tilde{g}_{ab})$ rather than in the physical one, $(\mathcal{M}, g_{ab})$. We now use this remark to put the results of Sec. 2 into a more general context extending the line of reasoning outlined in Sec. 3, and to establish some extra necessary and/or sufficient conditions for the different parts of $F_a^{(l)}$ to vanish.

Let us consider a charge with four-velocity v^a in an arbitrary conformally static⁸ spacetime $(\mathcal{M}, g_{ab})$. By definition, such a spacetime admits a hypersurface-orthogonal timelike conformal Killing vector field $\eta = \partial/\partial t$, corresponding to a conformal Killing time t . One can then define the scalar function

$$\Phi := \frac{1}{2} \ln(-g_{ab}\eta^a\eta^b) \quad (4.2)$$

and consider the ultrastatic spacetime $(\mathcal{M}, \tilde{g}_{ab})$ with $\tilde{g}_{ab}$ given by (4.1), in which η^a is a Killing vector field with unit norm, $\mathcal{L}_\eta \tilde{g}_{ab} = 0$ and $\tilde{g}_{ab}\eta^a\eta^b = -1$ [24]. It is not difficult to check that the projection, along the integral curves of η^a , of null geodesics of $(\mathcal{M}, g_{ab})$ onto the spatial hypersurfaces $\mathcal{S}$ defined by the condition $t = \text{const}$, are geodesics of $(\mathcal{S}, \tilde{h}_{ab})$, where $\tilde{h}_{ab} := \tilde{g}_{ab} + \eta_a\eta_b$ and $\eta_a := \tilde{g}_{ab}\eta^b$. Hence, generalising what we have done in Sec. 3, all quantities pertaining to the spacetime $(\mathcal{M}, \tilde{g}_{ab})$ will be denoted as “optical.” Instead of $F_a^{(1)}$, we can then consider $\tilde{F}_a^{(1)}$, computed in the optical spacetime $(\mathcal{M}, \tilde{g}_{ab})$ starting from the rescaled four-velocity $\tilde{v}^a = e^\Phi v^a$.

At any point on the particle worldline, $\tilde{v}^a$ can be decomposed in a unique way as

$$\tilde{v}^a = \gamma(\eta^a + v\tilde{\tau}^a), \quad (4.3)$$

where $v \in]-1, 1[$ is the speed of the particle according to an observer with four-velocity parallel to η^a , the coefficient $\gamma := (1 - v^2)^{-1/2}$ is the corresponding Lorentz factor, and $\tilde{\tau}^a$ is a spacelike vector orthogonal to η^a , such that $\tilde{g}_{ab}\tilde{\tau}^a\tilde{\tau}^b = 1$. The four-acceleration $\tilde{a}_a := \tilde{v}^b \tilde{\nabla}_b \tilde{v}_a$ of the particle in the optical spacetime $(\mathcal{M}, \tilde{g}_{ab})$ is then

$$\tilde{a}_a = \gamma^2 \tilde{T}_a \tilde{v}^b \tilde{\nabla}_b v + \gamma^2 v^2 \tilde{\tau}^b \tilde{\nabla}_b \tilde{\tau}_a + \gamma^2 v \eta^b \tilde{\nabla}_b \tilde{\tau}_a, \quad (4.4)$$

where

$$\tilde{T}^a := \frac{\tilde{k}^a{}_b \tilde{\tau}^b}{\sqrt{\tilde{k}_{cd} \tilde{\tau}^c \tilde{\tau}^d}} = \gamma(\tilde{\tau}^a + v\eta^a) \quad (4.5)$$

is a unit vector in $(\mathcal{M}, \tilde{g}_{ab})$, orthogonal to $\tilde{v}^a$. Using now the orthogonality condition $\eta^a \tilde{\tau}_a = 0$, and the property $\tilde{\nabla}_b \eta_a = 0$ that follows from the fact that $(\mathcal{M}, \tilde{g}_{ab})$ is ultrastatic, we have

$$\tilde{\nabla}_b \tilde{\tau}_a = \tilde{D}_b \tilde{\tau}_a - \eta_b \eta^c \tilde{\nabla}_c \tilde{\tau}_a, \quad (4.6)$$

where $\tilde{D}_b$ is the covariant derivative along directions orthogonal to η^a , hence on $(\mathcal{S}, \tilde{h}_{ab})$. In order for both Eqs. (4.4) and (4.6) to make sense, one needs to know $\tilde{\tau}^a$ off the particle worldline. The natural extension of $\tilde{\tau}^a$ is such that its integral curves (located on $t = \text{const}$ hypersurfaces) coincide with the projection on $\mathcal{S}$ of the particle worldline along the integral curves of η^a . This is equivalent to requiring that $\mathcal{L}_\eta \tilde{\tau}^a = 0$, i.e. that $\eta^b \tilde{\nabla}_b \tilde{\tau}^a = \tilde{\tau}^b \tilde{\nabla}_b \eta^a$, and the latter quantity vanishes in an ultrastatic spacetime. Hence, $\eta^b \tilde{\nabla}_b \tilde{\tau}_a = 0$, and the particle acceleration in optical spacetime takes the very simple form

$$\tilde{a}_a = \gamma^2 \tilde{T}_a \tilde{v}^b \tilde{\nabla}_b v + \gamma^2 v^2 \tilde{\tau}^b \tilde{D}_b \tilde{\tau}_a, \quad (4.7)$$

⁸This restriction is motivated mainly by simplicity. Optical geometry can be straightforwardly defined also for conformally stationary spacetimes [21], and has been extended even to cases where no global time symmetry exists, as it happens in gravitational collapse [22]. For recent results concerning generalisations of optical geometry, see Ref. [23].

in which one can identify a tangential and an orthogonal (centripetal) component [16].

The relationship [19]

$$\tilde{a}_a = a_a - k_a{}^b \nabla_b \Phi \quad (4.8)$$

generalises the result, already obtained in Sec. 3 for the special case of Schwarzschild spacetime, according to which one can write the particle acceleration in optical spacetime, $\tilde{a}_a$, as the sum between the non-gravitational thrust per unit mass, a_a , and a gravitational force per unit mass, $\tilde{g}_a = -k_a{}^b \nabla_b \Phi$:

$$\tilde{a}_a = a_a + \tilde{g}_a . \quad (4.9)$$

Equation (4.9) contains a precise formulation of the equivalence principle, because the quantity that is physically measurable at the particle location — the thrust, proportional to a_a — allows one only to determine the difference $\tilde{a}_a - \tilde{g}_a$ between $\tilde{a}_a$ and $\tilde{g}_a$, but not $\tilde{a}_a$ and $\tilde{g}_a$ separately (unless one possesses extra knowledge, of a non-local nature, about the structure of the spacetime under consideration).⁹ Inserting (4.7) into (4.9), and identifying $-\gamma^2 \tilde{T}_a \tilde{v}^b \tilde{\nabla}_b v$ and $-\gamma^2 v^2 \tilde{\tau}^b \tilde{D}_b \tilde{\tau}_a$ with suitable inertial forces per unit mass [16, 25], one finds a general relativistic version of the statement, commonly expressed within the framework of Newtonian mechanics, that in a frame comoving with the particle there is perfect balance between the non-gravitational thrust, the gravitational force, and the inertial forces acting on the particle.

On replacing (4.7) into the definition of $\tilde{F}_a^{(l)}$ one can find the local part of the self-force expressed in terms of v and $\tilde{\tau}^a$, that characterize the motion in optical spacetime $(\mathcal{M}, \tilde{g}_{ab})$. However, since the general expression is not particularly illuminating, let us focus instead on the most interesting particular cases.

4.1 Uniform motion along optical geodesics

It is obvious from Eq. (4.7) that a particle which moves uniformly (i.e., with $\tilde{v}^b \tilde{\nabla}_b v = 0$) along optical geodesics of space (so that $\tilde{\tau}^b \tilde{D}_b \tilde{\tau}_a = 0$) has vanishing acceleration in $(\mathcal{M}, \tilde{g}_{ab})$, that is, $\tilde{a}_a = 0$. Then $\tilde{F}_a^{(l)}$ consists only of the Hobbs term and we can write

$$F_a = e^{-\Phi} \left(\frac{1}{3} e^2 k_a{}^b \tilde{R}_{bc} \tilde{v}^c + e^2 \tilde{v}^b \int_{-\infty}^{\tilde{\tau}} d\tilde{\tau}' \tilde{f}_{aba'} \tilde{v}^{a'} \right) . \quad (4.10)$$

Thus, the particle is indeed subjected to a self-force, but just to the one that is associated with the geometric properties of the optical space. In a sense, the charge “feels” the geometry of $(\mathcal{M}, \tilde{g}_{ab})$ rather than the one of the physical spacetime $(\mathcal{M}, g_{ab})$.

It may be interesting to note that, with the only exception of ultrastatic spacetimes, for which the ordinary and the optical geometries simply coincide, there are no lines in the ordinary space $(\mathcal{S}, h_{ab})$ such that the von Laue term vanishes identically for a charge that moves uniformly along them. On the contrary, in $(\mathcal{S}, \tilde{h}_{ab})$ the conditions that $v = \text{const}$ and $\tilde{F}_a^{(vL)} = 0$ uniquely select optical geodesics.

⁹In particular, the two situations with $\tilde{a}_a = a_a$ and $\tilde{g}_a = 0$, or with $\tilde{a}_a = 0$ and $\tilde{g}_a = -a_a$, cannot be distinguished by a measurement of the thrust.

4.2 Special metrics

If $k_a{}^b \tilde{R}_{bc} \tilde{v}^c = 0$, then the expression (4.10) implies that $F_a = F_a^{(\text{nl})}$, because $F_a^{(1)}$ and $F_a^{(\text{nl})}$ do not mix under a conformal transformation [19].¹⁰ This can happen iff $\tilde{R}_a{}^b \tilde{v}_b = \lambda \tilde{v}_a$ for some λ , and it is easy to see, by contracting this equation with η^a and noting that $\eta^a \tilde{R}_a{}^b = 0$ in an ultrastatic spacetime, that it must be $\lambda = 0$. Using again the decomposition (4.3) and the property $\tilde{R}_{ab} \eta^b = 0$, one finally finds that the Hobbs term vanishes iff $v \tilde{R}_a{}^b \tilde{\tau}_b = 0$. Thus, excluding the trivial case $v = 0$, there is no local self-force for uniform motion along optical geodesics whose directions are eigenvectors of $\tilde{R}_a{}^b$ with zero eigenvalue. Such directions exist iff the determinant of the matrix made by the components $\tilde{R}_i{}^j$ with $i, j = 1, 2, 3$ is zero, i.e., iff such matrix is degenerate.

Let us reconsider the examples of Sec. 2 in the light of this conclusion. For the Einstein universe, $\tilde{R}_i{}^j = R_i{}^j = (2/R^2) \delta_i{}^j$, which is always non-degenerate. For Schwarzschild spacetime,

$$\tilde{R}_{\varphi i} = \frac{2M}{r} \left(1 - \frac{2M}{r}\right)^{-1} \left(1 - \frac{M}{r} (2 + \sin^2 \theta)\right) \delta_i{}^\varphi \quad (4.11)$$

and, for $\theta = \pi/2$, $\tilde{R}_i{}^j$ becomes degenerate at $r = 3M$. Hence, $F_a^{(1)}$ vanishes at $r = 3M$ for uniform motion in an equatorial plane along *any* optical geodesic, not only for the circular orbit considered in Sec. 2.2. This result is far from trivial, but its derivation is rather straightforward working in the optical space.

4.3 Geodesic motion

For charges following a geodesic in $(\mathcal{M}, g_{ab})$ the four-acceleration a_a vanishes, so the von Laue term $F_a^{(\text{vL})}$ is identically zero.¹¹ In optical geometry this result is not obvious, because it follows from a compensation between contributions coming from the “optical” von Laue and Hobbs terms. More precisely, applying (4.8) to this case one immediately finds $\tilde{a}_a = -k_a{}^b \nabla_b \Phi$. Then, the von Laue term $\tilde{F}_a^{(\text{vL})}$ in optical space contains first and second derivatives of Φ , that are canceled by identical contributions coming from the Hobbs term $\tilde{F}_a^{(\text{H})}$, because of the way the Ricci tensor changes under a conformal transformation [19].

Thus, from the perspective of optical geometry, the fact that $F_a^{(\text{vL})} = 0$ just when $\tilde{a}_a = -k_a{}^b \nabla_b \Phi$ is somewhat surprising. For example, it is not evident that $\tilde{F}_a^{(1)} = 0$ for the Keplerian motions of Sec. 2.2, because in optical geometry, where gravity is described by the physical field Φ , Keplerian motions have nothing qualitatively different from, say, orbits in a Coulomb field. In fact, it turns out that it is only thanks to a cancellation between the von Laue and the Hobbs terms in the optical geometry that $\tilde{F}_a^{(1)} = 0$. However, if one thinks that for a Keplerian motion in Schwarzschild spacetime $F_a^{(\text{vL})} = F_a^{(\text{H})} = 0$, and that $\tilde{F}_a^{(1)} = e^\Phi F_a^{(1)}$, then such a compensation is only expected. Geodesics in $(\mathcal{M}, g_{ab})$ are not geodesics in $(\mathcal{M}, \tilde{g}_{ab})$, so $\tilde{F}_a^{(\text{vL})}$ does not vanish in general, and since $\tilde{R}_{ab} \neq 0$, the Hobbs force $\tilde{F}_a^{(\text{H})}$ also does not vanish. Nevertheless, $\tilde{F}_a^{(\text{vL})}$ and $\tilde{F}_a^{(\text{H})}$ combine to form a vanishing local self-force, $\tilde{F}_a^{(1)} = 0$.

¹⁰For a more general, non-uniform motion, one finds $\tilde{F}_a^{(1)} = \tilde{F}_a^{(\text{vL})}$.

¹¹We avoid referring to this situation as “free fall,” because the latter would be appropriately identified by the vanishing of the external force in Eq. (1.1), $K_a = 0$, whereas geodesic motion corresponds to $K_a + F_a = 0$.

This remark should not be regarded as a drawback of the description based on optical geometry, though. Indeed, the fact that $F_a^{(\text{vL})}$ does, or does not, vanish is not particularly important, because the physically interesting quantity is not $F_a^{(\text{vL})}$, but $F_a^{(\text{l})}$. And whether $F_a^{(\text{l})} = 0$ (or, equivalently, $\tilde{F}_a^{(\text{l})} = 0$) cannot be established simply by looking at $F_a^{(\text{vL})}$ (or $\tilde{F}_a^{(\text{vL})}$), unless one is in a situation for which $F_a^{(\text{H})} = 0$ (or $\tilde{F}_a^{(\text{H})} = 0$). Thus, in the case of Schwarzschild spacetime the evaluation of $F_a^{(\text{vL})}$ turns out to be convenient only because, since $R_{ab} = 0$ there, the Hobbs force vanishes identically, so $F_a^{(\text{vL})}$ coincides with $F_a^{(\text{l})}$. However, in a spacetime for which it is $\tilde{R}_{ab}$ that vanishes, one has $\tilde{F}_a^{(\text{H})} = 0$. Consequently, it is now $\tilde{F}_a^{(\text{vL})}$ that one should inspect, in order to get information about the behaviour of $F_a^{(\text{l})} = 0$.

4.4 Conformally static charge

In order to further illustrate the point made in the last paragraph of Sec. 4.3, let us consider a conformally static charge, i.e., one with four-velocity $v^a = n^a$, where

$$n^a := e^{-\Phi} \eta^a = \left(-g_{bc} \eta^b \eta^c \right)^{-1/2} \eta^a \quad (4.12)$$

is the unit vector field parallel to η^a . This is obviously a subcase of the situations covered in Sec. 4.1, and since $\tilde{R}_{ab} \eta^b = 0$, it is evident in optical geometry that $\tilde{F}_a^{(\text{vL})} = \tilde{F}_a^{(\text{H})} = 0$, so also $\tilde{F}_a^{(\text{l})} = 0$ and $F_a^{(\text{l})} = 0$. On the other hand, this conclusion is not obvious at all if one works in the spacetime $(\mathcal{M}, g_{ab})$, because in order to establish it, one must use highly non-trivial properties of the vector field n^a , that lead to cancelations between terms coming from $F_a^{(\text{vL})}$ and $F_a^{(\text{H})}$ (none of which, however, vanishes separately). Hence, in this case it is optical geometry that allows one to establish that $F_a^{(\text{l})} = 0$ in a simple way, by using the decomposition $\tilde{F}_a^{(\text{l})} = \tilde{F}_a^{(\text{vL})} + \tilde{F}_a^{(\text{H})}$. The alternative split, $F_a^{(\text{l})} = F_a^{(\text{vL})} + F_a^{(\text{H})}$, leads instead to rather cumbersome calculations.

5 Conclusions

Let us summarise the results of Secs. 2–4. We have seen that, for uniform motion along a spatial geodesic $\chi = \pi/2$ in Einstein's static universe, the von Laue force vanishes, whereas the Hobbs force does not (hence, the local part of the self-force also does not vanish). In Schwarzschild spacetime, for uniform motion along a circle at $r = 3M$, the von Laue force vanishes; since the Hobbs force is identically zero (Schwarzschild spacetime is Ricci-flat), this implies that $F_a^{(\text{l})}(3M) = 0$. Similarly, in the optical Schwarzschild spacetime, $\tilde{F}_a^{(\text{vL})}(3M) = \tilde{F}_a^{(\text{H})}(3M) = \tilde{F}_a^{(\text{l})}(3M) = 0$. All these are instances of the following general results, valid in an arbitrary conformally static spacetime: (i) for geodesic motion in optical spacetime $(\mathcal{M}, \tilde{g}_{ab})$ — uniform motion along optical geodesics —, $\tilde{F}_a^{(\text{vL})} = 0$; (ii) for geodesic motion in the spacetime $(\mathcal{M}, g_{ab})$, $F_a^{(\text{vL})} = 0$. Note that, however, the physically interesting result does not concern $F_a^{(\text{vL})}$ or $\tilde{F}_a^{(\text{vL})}$, but the entire local part of the self-force. Whether this vanishes in cases (i) and (ii) depends on the additional feature that the matrix $\tilde{R}_i^j$ or R_μ^ν , respectively, be degenerate. Also, note that the particular case of (i), that for a conformally static charge $F_a^{(\text{l})} = \tilde{F}_a^{(\text{l})} = 0$, although non-trivial, is straightforward when regarded from the perspective offered by optical geometry. Hence, the latter appears to be a useful tool in calculations of self-force and related effects, as it happens already in several other circumstances [15, 16, 17, 20, 21, 22, 25, 26, 27, 28].

Indeed, although in this paper we deliberately avoided dealing with the non-local part of the self-force $F_a^{(\text{nl})}$, there are good reasons to believe that working in the optical spacetime might also simplify its evaluation — a rather challenging task, in general. This is suggested by the fact that computing $F_a^{(\text{nl})}$ amounts, basically, to finding the bi-tensor $f_{aba'}$ in Eq. (1.3). In turn, this amounts to the determination of a particular electromagnetic field in spacetime, and we know already from Ref. [28] that this is sometimes easier to do using optical geometry. We leave the development of this subject for future investigations.

Acknowledgements

It is a pleasure to thank Rossella Rosin for her careful preparation of the figures. S.S. is grateful to Nordita and to the Department of Astronomy and Astrophysics at Chalmers University for hospitality.

Appendix: Centripetal acceleration in optical geometry

We present a derivation of Eq. (3.25) based on the embedding diagram of Fig. 2. In the fictitious three-dimensional Euclidean space, the centripetal acceleration of a particle that moves uniformly on the orbit $r = \text{const}$ is $a_E^{(\text{c})} = \gamma^2 \Omega^2 \rho$, where $\rho = (1 - 2M/r)^{-1/2} r$ is the distance from the axis [18] — also called “radius of gyration” [26] because $v = (1 - 2M/r)^{-1/2} \Omega r = \Omega \rho$. The centripetal acceleration in the optical space is given by the tangential component of a_E , namely

$$a^{(\text{c})} = \gamma^2 \Omega^2 \rho \cos \alpha, \quad (\text{A1})$$

where α is the angle indicated in Fig. 2. Let $z = f(\rho)$ be the equation of the surface in Fig. 2. The surface is isometric to the section $\theta = \pi/2$ of the optical space if [18]

$$\left(1 + \left(\frac{df}{d\rho}\right)^2\right) d\rho^2 = \left(1 - \frac{2M}{r}\right)^{-2} dr^2. \quad (\text{A2})$$

But we have also $df/d\rho = \tan \alpha$, so

$$\cos \alpha = \left(1 - \frac{2M}{r}\right) \left|\frac{d\rho}{dr}\right| = \left(1 - \frac{2M}{r}\right)^{-1/2} \left|1 - \frac{3M}{r}\right|. \quad (\text{A3})$$

Substituting into (A1) we recover Eq. (3.25).

References

- [1] R. M. Wald, *General Relativity* (University of Chicago Press, Chicago, 1984).
- [2] B. S. DeWitt and R. W. Brehme, “Radiation damping in a gravitational field,” *Ann. Phys.* (N.Y.) **9**, 220–259 (1960).

- [3] J. M. Hobbs, “A vierbein formalism of radiation damping,” *Ann. Phys. (N.Y.)* **47**, 141–165 (1968).
- [4] F. Rohrlich, *Classical Charged Particles* (Addison-Wesley, Reading, 1990).
- [5] F. Rohrlich, “The electron: development of the first elementary particle theory,” in *The Physicist’s Conception of Nature*, edited by J. Mehra (Reidel, Dordrecht, 1973), pp. 331–369;
 ——— “The dynamics of a charged sphere and the electron,” *Am. J. Phys.* **65**, 1051–1056 (1997).
- [6] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*, 4th edition (Butterworth-Heinemann, Oxford, 1975), pp. 224–225, 227.
 T. C. Quinn and R. M. Wald, “Axiomatic approach to electromagnetic and gravitational radiation reaction of particles in curved spacetime,” *Phys. Rev. D* **56**, 3381–3394 (1997) [gr-qc/9610053].
- [7] S. Sonego and V. Faraoni, “Huygens’ principle and characteristic propagation property for waves in curved space-times,” *J. Math. Phys.* **33**, 625–632 (1992).
 V. Faraoni and S. Sonego, “On the tail problem in cosmology,” *Phys. Lett. A* **170**, 413–420 (1992) [astro-ph/9209004].
 L. Bombelli and S. Sonego, “Relationships between various characterizations of wave tails,” *J. Phys. A* **27**, 7177–7199 (1994) [math-ph/0002026].
 B. C. Nolan, “A characterization of strong wave tails in curved spacetimes,” *Class. Quantum Grav.* **14**, 1295–1308 (1997) [gr-qc/9703031].
- [8] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, Cambridge, 1973), p. 80.
- [9] A. D. Yaghjian, *Relativistic Dynamics of a Charged Sphere* (Springer, Berlin, 1992).
 F. Rohrlich, “Classical self-force,” *Phys. Rev. D* **60**, 084017 (1999).
 S. Parrott, “Asymptotics of a proposed delay-differential equation of motion for charged particles,” e-print gr-qc/0205065;
 ——— “Comment on *Phys. Rev. D* **60** 084017 “Classical self-force” by F. Rohrlich,” e-print gr-qc/0502029.
 A. I. Harte, “Self-forces on extended bodies in electrodynamics,” *Phys. Rev. D*, to appear [gr-qc/0508123].
- [10] E. Poisson, “The motion of point particles in curved spacetime,” *Living Rev. Relativity* **7**, 6 (2004); URL (cited on February 16, 2006): <http://www.livingreviews.org/lrr-2004-6> [gr-qc/0306052].
 S. Detweiler, “Perspective on gravitational self-force analysis,” *Class. Quantum Grav.* **22**, S681–S716 (2005) [gr-qc/0501004].
- [11] S. Weinberg, *Gravitation and Cosmology* (Wiley, New York, 1972), p. 471.
- [12] L. M. Burko, “Self-force approach to synchrotron radiation,” *Am. J. Phys.* **68**, 456–468 (2000) [gr-qc/9902079].

- [13] M. A. Abramowicz and J. P. Lasota, “A note on a paradoxical property of the Schwarzschild solution,” *Acta Phys. Pol.* **B5**, 327–329 (1974);
 ——— “On traveling round without feeling it and uncurving curves,” *Am. J. Phys.* **54**, 936–939 (1986).
- [14] A. A. Sokolov, D. V. Galtsov, and V. I. Petukhov, “Radiation emitted by relativistic particles moving in the vicinity of the Schwarzschild black hole, immersed in an external magnetic field,” *Phys. Lett. A* **68**, 1–2 (1978).
- [15] J. S. Dowker and G. Kennedy, “Finite temperature and boundary effects in static spacetimes,” *J. Phys. A* **11**, 895–920 (1978).
 G. W. Gibbons and M. J. Perry, “Black holes and thermal Green functions,” *Proc. R. Soc. Lond. A* **358**, 467–494 (1978).
 M. A. Abramowicz, B. Carter, and J. P. Lasota, “Optical reference geometry for stationary and static dynamics,” *Gen. Relativ. Gravit.* **20**, 1173–1183 (1988).
- [16] M. A. Abramowicz, “Inertial forces in general relativity,” in *The Renaissance of General Relativity and Cosmology*, edited by G. Ellis, A. Lanza, and J. Miller (Cambridge University Press, Cambridge, 1993), pp. 40–58.
 M. A. Abramowicz, P. Nurowski, and N. Wex, “Covariant definition of inertial forces,” *Class. Quantum Grav.* **10**, L183–L186 (1993).
 S. Sonego and M. Massar, “Covariant definition of inertial forces: Newtonian limit and time-dependent gravitational fields,” *ibid.* **13**, 139–144 (1996).
- [17] M. A. Abramowicz, “Centrifugal force: a few surprises,” *Mon. Not. R. Astron. Soc.* **245**, 733–746 (1990);
 ——— “Relativity of inwards and outwards: an example,” *ibid.* **256**, 710–718 (1992).
 M. A. Abramowicz and E. Szuszkiewicz, “The wall of death,” *Am. J. Phys.* **61**, 982–991 (1993).
- [18] S. Kristiansson, S. Sonego, and M. A. Abramowicz, “Optical space of the Reissner–Nordström solutions,” *Gen. Relativ. Gravit.* **30**, 275–288 (1998).
- [19] S. Sonego, “Conformal behavior of the Lorentz-Dirac equation and Machian particle dynamics,” *J. Math. Phys.* **40**, 3918–3924 (1999).
- [20] M. A. Abramowicz and A. R. Prasanna, “Centrifugal-force reversal near a Schwarzschild black hole,” *Mon. Not. R. Astron. Soc.* **245**, 720–728 (1990).
 M. A. Abramowicz and J. C. Miller, “Ellipticity behaviour of relativistic Maclaurin spheroids,” *ibid.* **245**, 729–732 (1990).
 M. A. Abramowicz and J. Bičák, “On the interplay between relativistic gravitational, centrifugal and electric forces: a simple example,” *Gen. Relativ. Gravit.* **23**, 941–946 (1991).
 S. Sonego and M. Massar, “On the notions of gravitational and centrifugal force in static spherically symmetric spacetimes,” *Mon. Not. R. Astron. Soc.* **281**, 659–665 (1996).
 M. A. Abramowicz, “Rayleigh and Solberg criteria reversal near black holes: the optical geometry explanation,” e-print astro-ph/0411718.
- [21] G. F. Torres del Castillo and J. Mercado-Pérez, “Three-dimensional formulation of the Maxwell equations for stationary spacetimes,” *J. Math. Phys.* **40**, 2882–2890 (1999).

- [22] S. Sonego, J. Almergren, and M. A. Abramowicz, “Optical geometry for gravitational collapse and Hawking radiation,” *Phys. Rev. D* **62**, 064010 (2000) [gr-qc/0005106].
- [23] R. Jonsson and H. Westman, “Generalizing optical geometry,” *Class. Quantum Grav.* **23**, 61–76 (2006) [gr-qc/0403004].
- [24] S. A. Fulling, “Alternative vacuum states in static space-times with horizons,” *J. Phys. A* **10**, 917–951 (1977).
- [25] W. Hasse and V. Perlick, “Gravitational lensing in spherically symmetric static spacetimes with centrifugal force reversal,” *Gen. Relativ. Gravit.* **34**, 415–433 (2002) [gr-qc/0108002].
T. Foertsch, W. Hasse, and V. Perlick, “Inertial forces and photon surfaces in arbitrary spacetimes,” *Class. Quantum Grav.* **20**, 4635–4651 (2003) [gr-qc/0306042].
W. Hasse and V. Perlick, “A Morse-theoretical analysis of gravitational lensing by a Kerr-Newman black hole,” e-print gr-qc/0511135.
- [26] M. A. Abramowicz, J. C. Miller, and Z. Stuchlík, “Concept of radius of gyration in general relativity,” *Phys. Rev. D* **47**, 1440–1447 (1993).
- [27] S. Sonego and A. Lanza, “Relativistic perihelion advance as a centrifugal effect,” *Mon. Not. R. Astron. Soc.* **279**, L65–L66 (1996).
M. A. Abramowicz, A. Lanza, J. C. Miller, and S. Sonego, “Curving Newtonian space,” *Gen. Relativ. Gravit.* **29**, 1585–1596 (1997).
M. A. Abramowicz, N. Andersson, M. Bruni, P. Ghosh, and S. Sonego, “Gravitational waves from ultracompact stars: the optical geometry view of trapped modes,” *Class. Quantum Grav.* **14**, L189–L194 (1997).
M. A. Abramowicz and W. Kluźniak, “Epicyclic orbital oscillations in Newton’s and Einstein’s dynamics,” *Gen. Relativ. Gravit.* **35**, 69–77 (2003) [gr-qc/0206063].
- [28] S. Sonego and M. A. Abramowicz, “Maxwell equations and the optical geometry,” *J. Math. Phys.* **39**, 3158–3166 (1998).